

Noise-assisted preparation of entangled atoms

X. X. Yi, C. S. Yu, L. Zhou, and H. S. Song

Department of physics, Dalian University of Technology, Dalian 116024, China

(Dated: February 1, 2008)

We discuss the generation of entangled states of two two-level atoms inside an optical cavity. The cavity mode is supposed to be coupled to a white noise with adjustable intensity. We describe how the entanglement between the atoms inside the cavity arise in such a situation. The entanglement is maximized for intermediate values of the noise intensity, while it is a monotonic function of the spontaneous rate. This resembles the phenomenon of stochastic resonance and sheds more light on the idea to exploit white noise in quantum information processing.

PACS numbers: 03.67.-a, 03.67.-Hz

Entanglement shared between distant sites is a valuable resource for quantum information processing [1, 2, 3]. Stimulated by this discovery, there are a lot of attention have been focused on how to create, manipulate and exploit entanglement from both sides of experimental and theoretical studies [4]. From a practical point of view, the main task to create and exploit entanglement is how to minimize the impact of noise that is not able to be isolated from our system in any real experimental scenario. This interaction between the system of interest and the noise which usually models surroundings of the system results in a decoherence process. As a consequence the entangled system may end up in a mixed state that would be no longer useful for any quantum information processing. It is therefore important for practical realization of quantum information processing protocols to engineer mechanisms to prevent, minimize, or use the impact of environmental noise.

Numerous proposals have been made for preventing, minimizing or using the environmental noise, for example, loop control strategies, that use an ancillary system coupling to the quantum processor to better the performance of the proposals[5, 6], quantum error correction [7] uses redundant coding to protect quantum states against noisy environments. This procedure is successful as long as the error rate is sufficiently small. It wastes a number of qubits and quantum gates, and then limit its implementation by present available technology. A more economic approach consists of exploiting the existence of so-called decoherence-free subspace that are completely insensitive to specific types of noise [8]. This approach tends to require fewer additional resources, but is only applicable in specific situations. The seminal idea that dissipation can assist the generation of entanglement has been put forward recently [9, 10, 11]. In a system consisting of two distinct leaky optical cavities, it was shown that the entanglement is maximized for intermediate values of the cavity damping rates and the intensity of the white noise, vanishing both for small and for large values of these parameters [11]. In fact, this idea appeared first in Ref.[9] for two atoms inside an optical cavity and it shows that cavity decay can assist the preparation of




FIG. 1: Schematic diagram for the preparation of entangled two atoms. The atom-cavity system is prepared initially in the ground state $|g\rangle_a|g\rangle_b$ with no photon populated in the cavity. The atoms become entangled and the entanglement is maximized for intermediate values of the noise intensity.

maximally entangled atoms, without cavity decay, the reduced state of the two-atom system would be in an inseparable mixture at all times, but not in a maximally entangled one.

In the latter case, the photon leakage leads to the undesired parts of the global wave-function to decay and therefore such terms are eliminated for sufficiently large times. This requires that the overlapping between the initial state and the desired maximally entangled state must not be zero, hence one of the two atoms prepared in its excited state initially is necessary. This point makes our proposed scheme here for preparation of entangled two atoms different from the above one.

In this letter, we put the idea in Ref. [11] forward to a two-atom system, we use noise to play a constructive role in quantum information processing. We will concentrate on the problem of creating entangled atoms when only incoherent sources are available and demonstrate that, indeed, controllable entanglement can arise in this situation.

Our system consists of two two-level atoms inside a leaky optical cavity. As depicted in figure 1, we will re-

fer to atom a and atom b when the context requires us to differentiate them, but otherwise they are supposed to be identical. We denote the atomic ground and excited states by $|g\rangle_i$ and $|e\rangle_i$, respectively, and call 2Γ ($\Gamma_a = \Gamma_b = \Gamma$) the spontaneous emission rate from the upper level. We assume that the distance between the atoms is much larger than an optical wavelength, therefore dipole-dipole interaction can be neglected. The cavity mode is assumed to be resonant with the atomic transition frequency, and we will denote the cavity decay rate by 2κ . For the sake of generality we allow the coupling between each atom and the cavity mode, g_i , to be different. We suppose the cavity is driven by an external thermal field (white noise) whose intensity will be characterized in terms of an effective photon number n_T , but we do not want to specify the noise. The relaxation of the atom-cavity system can take place through two channels, at rate 2κ (cavity decay) and 2Γ (spontaneous decay). The master equation governing the time evolution of the global system is given by (setting $\hbar = 1$)

$$\dot{\rho} = -i[H, \rho] + \mathcal{L}(\rho), \quad (1)$$

where the Hamiltonian H describes the internal energies of the cavity and the two atoms as well as the atom-cavity coupling. The Liouvillian $\mathcal{L}(\rho)$ describes the atom decay and the interaction of the cavity mode with the noise. As no external coherent driving is present, the Hamiltonian reads

$$H = \frac{\omega}{2}\sigma_a^z + \frac{\omega}{2}\sigma_b^z + \omega_f a^\dagger a + \sum_{i=a,b} g_i(|g\rangle_i\langle e|a^\dagger + h.c.), \quad (2)$$

where a represent the cavity mode with frequency ω_f . The Liouvillian is given by

$$\begin{aligned} \mathcal{L}(\rho) = & -\kappa(n_T + 1)(a^\dagger a \rho + \rho a^\dagger a - 2a \rho a^\dagger) \\ & - \kappa n_T (a a^\dagger \rho + \rho a a^\dagger - 2a^\dagger \rho a) \\ & - \Gamma \sum_{i=a,b} (|e\rangle_i\langle e| \rho + \rho |e\rangle_i\langle e| - 2|g\rangle_i\langle e| \rho |e\rangle_i\langle g|) \end{aligned} \quad (3)$$

Here Γ describes the atom decay rate and κ stands for the cavity leakage rate. We do not explicitly specify the white noise, but its intensity n_T refers to its effective particle number. To simplify the representation, now we turn to an interaction picture with respect to $H_0 = \sum_{i=a,b} \frac{\omega}{2}\sigma_i^z + \omega a^\dagger a$. After this transformation, the Liouvillian part remains unchanged, while the Hamiltonian part is now given by

$$H_I = \sum_{i=a,b} g_i(|g\rangle_i\langle e|a^\dagger + h.c.), \quad (4)$$

where that the atom-cavity coupling is on resonance was assumed. The analytical solution to the equation (1) is extremely tedious. To make the physical interpretation clear, we introduce two new effective atomic modes one

of which will be decoupled from the cavity mode. The two collective atomic modes is given by the following definition

$$\sigma_A^+ = \frac{g_a \sigma_a^+ + g_b \sigma_b^+}{\sqrt{g_a^2 + g_b^2}}, \sigma_B^+ = \frac{g_b \sigma_a^+ - g_a \sigma_b^+}{\sqrt{g_a^2 + g_b^2}}, \quad (5)$$

with $\sigma_i^+ = |e\rangle_i\langle g|$, together with $\sigma_i^- = (\sigma_i^+)^\dagger$, and σ_i^z represent the pauli operators for the atom i . In terms of these new operators, the Hamiltonian and Liouvillian part of the master equation are given by

$$H_I = g(|g\rangle_A\langle e|a^\dagger + h.c.), \quad (6)$$

where $g = \sqrt{g_a^2 + g_b^2}$, and

$$\begin{aligned} \mathcal{L}(\rho) = & -\kappa(n_T + 1)(a^\dagger a \rho + \rho a^\dagger a - 2a \rho a^\dagger) \\ & - \kappa n_T (a a^\dagger \rho + \rho a a^\dagger - 2a^\dagger \rho a) \\ & - \Gamma \sum_{i=A,B} (|e\rangle_i\langle e| \rho + \rho |e\rangle_i\langle e| - 2|g\rangle_i\langle e| \rho |e\rangle_i\langle g|) \end{aligned} \quad (7)$$

Note that the sum in the last line of Eq.(7) is taken over the two NEW modes. The transformation between the resulting atom a, b and the collective modes A, B is clear. For example, both the resulting atoms a and b in its ground state $|g\rangle_a|g\rangle_b$ can be equivalently expressed in terms of $|g\rangle_A|g\rangle_B$ and $|e\rangle_a|e\rangle_b$ likewise. The new master equation Eq.(7) shows us that we have one mode (mode B) which is completely decoupled from the Hamiltonian dynamics and is purely damped under the Liouvillian dynamics, this is a consequence of the transformation from the resulting atoms to the collective modes. The mode B then will not be populated in steady state irrespective of its initial states. In other words, if the mode B is in its ground state initially, it will remain on that forever. Therefore, we begin our investigations with both collective modes A and B in the ground state $|g\rangle_A|g\rangle_B$. As the mode B will then never be populated, we disregard that mode in the following discussions. Apart from the above assumption, we discuss the entanglement generation here only for the case of no photon in the cavity initially, this is relevant to the topics under our consideration, i.e., study the role of the white noise in the entanglement generation. To understand the origin of the generation of entanglement from white noise, let us begin by considering a special case of perfect cavity, i.e., $\kappa = 0$. For the general form of the reduced density matrix of the atom system

$$\rho_{atom} = \rho_{ee}|e\rangle\langle e| + \rho_{eg}|e\rangle\langle g| + \rho_{ge}|g\rangle\langle e| + \rho_{gg}|g\rangle\langle g|,$$

it is easy to check that $\rho_{eg} = \rho_{ge} = 0$ in this situation, where $\rho_{xy}(x, y = g, e) = \text{Tr}_c(|y\rangle\langle x| \rho)$ and Tr_c denotes trace over the cavity mode. Noticing $|e\rangle_A\langle e| = \frac{g_a^2|e\rangle_a\langle e| + g_b^2|e\rangle_b\langle e|}{g_a^2 + g_b^2}$, and $|g\rangle_A\langle g|$ has a similar form, we conclude that for perfect cavity with no initial photon inside the cavity, the atoms remain separable at all the




FIG. 2: Wootters concurrence of the two-atom system as a function of the intensity of the white noise n_T and time t . The chosen parameters are $\omega = \omega_f$, $g_a = g_b = 1$, $\kappa = 2$ and $\Gamma = 0.2$. Note that the entanglement arrives at its maximum for an intermediate noise intensity.

times. Moreover this conclusion holds even for $\kappa \neq 0$ with $n_T = 0$. In this case, the cavity mode will never be populated if there is no photon in the cavity initially. As the cavity mode act as a data bus, no photon exist inside the cavity means the data bus is idle. This conclusion will be changed if we increase the value of the thermal noise intensity n_T . For a limiting case of $g = 0$ and $\Gamma = 0$, the stationary state of the cavity field is no longer the vacuum, instead, it takes[12]

$$\rho_c = \sum_n |n\rangle\langle n| \left(\frac{n_T}{1+n_T} \right)^n \frac{1}{1+n_T}.$$

The active data bus indicate that the two atoms inside the cavity will be entangled together.

We will now show in a numerical way that the envisioned idea we have described above can be realized in the atom-cavity system by properly chosen parameters. We will choose the Wootters concurrence as the entanglement measure [13],

$$c(\rho) = \max\{0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4\},$$

where the λ_i are the square roots of the eigenvalues of the non-Hermitian matrix $\rho\tilde{\rho}$ with $\tilde{\rho} = (\sigma_y \otimes \sigma_y)\rho^*(\sigma_y \otimes \sigma_y)$ in decreasing order. The Wootters concurrence gives an explicit expression for the entanglement of formation, which quantifies the resources needed to create a given entangled state. The typical behavior of the entanglement in the system is illustrated in figure 2. There we have plotted the amount of entanglement of the joint state of the two atoms as a two-variable function of the intensity of the noise n_T and time t . The chosen parameters are $\omega = \omega_f$, $g_a = g_b = 1$, $\kappa = 2$ and $\Gamma = 0.2$. We want to stress that our simulation is presented for Eq.(1), i.e., the original master equation for the atom-cavity system, and as we mentioned above the initial state of the global system is $|g\rangle_a|g\rangle_b|0\rangle_c$ in our simulation, and we cut off the




FIG. 3: The dependence of amount of entanglement on the noise intensity n_T and the cavity decay rate κ for a value of time $t = 1/(2g)$. The parameters chosen are $\omega = \omega_f$, $g_a = g_b = 1$, and $\Gamma = 0.2$. The system exhibits a resonance for both n_T and κ at an intermediate value.

intra-cavity photon number at a value of 5. Note that for any value of t in the region of entanglement $\neq 0$, the behavior of the amount of entanglement between the two atoms is non-monotonic, it increase to a maximum value for an optimal intensity of the noise and then decrease towards zero for a sufficiently large intensity. Physically, to get non-zero amount of entanglement, the cavity mode must be populated (or be excited) at any value of time t . For the two limiting case of either $n_T = 0$ or $n_T \rightarrow \infty$, however, the data bus remains idle for all the times. Thus the amount of entanglement equals zero. It is also worthwhile to study the dependence of entanglement on both the intensity of the noise and the cavity decay rates. In figure 3, we present those for $\omega = \omega_f$, $g_a = g_b = 1$, and $\Gamma = 0.2$. It seems that the amount of entanglement is fixed for a specific value of $n_T\kappa$. It is interesting to note that the amount of entanglement behave as a monotonic function of the atom decay rate Γ . As figure 4 shows, this is quite different from the case presented in Ref. [11], where the cavity decay can assist themselves to entangle together, and the entanglement is generated among themselves then.

The noise-assisted entanglement preparation is somehow reminiscent of the well known phenomenon of stochastic resonance [14, 15, 16], where the response of a system to a periodic force can be enhanced in the presence of an intermediate amount of noise. A related effect that cavity decay can assist the generation of squeezing has been found recently [17], there they show that the squeezing effect is enhanced as the damping rate of the cavity is increased to some extent. However, the pumping field amplitude is required to be inversely proportional to the damping rate for the optimal squeezing. This is similar to our results shown in figure 3.

To sum up, we have described an experimental situation where entanglement between two atomic




FIG. 4: Amount of entanglement versus the noise intensity n_T and the atomic spontaneous emission rate Γ for a value of time $t = 1/(2g)$. We choose the parameters $g_a = g_b = 1$, $\omega_f = \omega$ and $\kappa = 2$ for this plot. The entanglement of the two atoms is a monotonic function of Γ for any value of n_T , this is quite different from the case presented in Ref.[11]

systems can be prepared with assistance of the white noise. The entanglement measured by the Wootters concurrence is maximized for intermediate values of the cavity decay rate and the intensity of the white noise, while it is a monotonic function of the atomic spontaneous emission rate. Recall that the atomic decay itself can not induce entanglement among the atoms, even if at finite temperatures, we conclude that the coupling between the data bus and the white noise is the origin of the generation of the entanglement. The phenomenon of white noise-assisted entanglement generation is not a rare phenomenon, it resembles the phenomenon of stochastic resonance. However, this discovery [11] is really valuable because it sheds new light on the constructive role that noise may play in quantum information processing. In contrast with the results in Ref.[11], the proposal presented here is for the entanglement generation between two two-level atoms. For such a two-qubit system, any amount of entanglement, even if very small, is distillable [18], and therefore the entangled atoms are useful for quantum information processing. The entanglement among many atoms also can be created by the same manner, the

results will be presented elsewhere.

ACKNOWLEDGEMENT:

This work is supported by EYTP of M.O.E, and NSF of China.

-
- [1] C. H. Bennett and D. P. DiVincenzo, *Nature* **404**, 247(2000).
 - [2] M. A. Nielsen and I. L. Chuang, *Quantum computation and quantum information* (Cambridge University press, Cambridge, 2000).
 - [3] M. B. Plenio and V. Vedral, *Contemp. Phys.* **39**,431(1998).
 - [4] See, for example, J. M. Raimond *et al.*, *Rev. Mod. Phys.* **73**, 565(2001); Vedral, *Rev. Mod. Phys.* **74**, 197(2002).
 - [5] H. M. Wiseman and G. J. Milburn, *Phys. Rev. Lett.* **70**,548(1993); D. Vital, *et al.*, *Phys. Rev. Lett.* **79**, 2442(1997); L. Viola,*et al.*, *Phys. Rev. Lett.* **82**, 2417(1999).
 - [6] S. Mancini, *et al.*, *EuroPhys. Lett.* **60**, 498(2002).
 - [7] P. W. Shor, *Phys. Rev. A* **52**, 2493(1995); A. R. Calderbank and P. W. Shor, *Phys. Rev. A* **54**, 1098(1996); A. M. Steane, *Proc. Roy. Soc. A* **452**, 2551(1996).
 - [8] G. M. Palma, *et al.*, *Proc. Roy. Soc. A* **452**, 567(1996); M. B. Plenio, *et al.*, *Phys. Rev. A* **55**, 67(1997); D. A. Lidar, *et al.*, *Phys. Rev. Lett.* **81**, 2594(1998); A. Beige, *et al.*, *Phys. Rev. Lett.* **85**, 1762(2000); A. Beige, *et al.*, *New J. Phys.* **2**, 22(2000).
 - [9] M. B. Plenio, *et al.*, *Phys. Rev. A* **59**, 2468(1999); A. Beige, *et al.*, *J. Mod. Opt.* **47**, 2583(2000); P. Horodecki, *Phys. Rev. A* **63**,022108(2001).
 - [10] S. Bose, *et al.*, *Phys. Rev. Lett.* **83**, 5158(1999).
 - [11] M. B. Plenio, *et al.*, *Phys. Rev. Lett.* **88**, 197901(2002).
 - [12] C. W. Gardiner and P. Zoller, *Quantum Noise* (Springer 2000).
 - [13] W. K. Wootters, *Phys. Rev. Lett.* **80**, 2245(1998).
 - [14] L. Gamaitoni, *et al.*, *Rev. Mod. Phys.* **70**, 223(1998).
 - [15] S. F. Huelga and M. B. Plenio, *Phys. Rev. A* **62**, 052111(2000).
 - [16] A. Buchleitner, and R. N. Mantegna, *Phys. Rev. Lett.* **80**, 3932(1998).
 - [17] H. Nha, *et al.*, *quant-ph/0106054*.
 - [18] M. Horodecki, *et al.*, *Phys. Rev. A* **59**, 4206 (1999); X. X. Yi,*et al.*, *Phys. Rev. A* **62**, 062312(2000).